



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Time or Expence to make, and therefore might be supplied by new ones, as often as they happen'd to break.

These Moulds seem to have been burnt or baked sufficiently to make them hard, but not so as to render them porous like Bricks, whereby they would have lost their smooth and even Surface; which in these is plainly so close, that whatever Metal should be formed in them would have no Appearances like the Sand-Holes, by which counterfeit Coins or Medals are usually detected.

London, May 20.

1747.

XXVI. *An extract of a letter from William Jones Esq; F. R. S. to Martin Folkes Esq; President of the Royal Society; containing a commodious disposition of equations for exhibiting the relations of goniometrical lines.*

T H E O R E M.

Presented July 4. **I***N a circle whose radius is r, let there*
 1747. *be two arcs, A the greater, a the less, each in the first quadrant; put s, t, s', and v', for the sine, tangent, secant, and versed sine of an arc; s', t', t', the sine, tangent, secant of the complement, and v', the versed sine of the supplement of that arc; let $z = \frac{1}{2}A + a$, $x = \frac{1}{2}A - a$; or if z and x be put for the arcs, it will be $A = z + x$, $a = z - x$.*

Then will the terms in any column of the following table, be proportional to their corresponding ones in any other column.

From

*A TABLE of the Relations
of Goniometrical Lines.*

$z s, z$	$s, z z$ $s, A + a$	$u, z z$ $u, A + a$	$s, A + s, a$	$s, a - s, A$ $u, A - u, a$
$z s, x$	$s, A - s, a$	$s, a - s, A$ $u, A - u, a$	$s, z x$ $s, A - a$	$u, z x$ $u, A - a$
$z s', z$	$s', z z$ $s', A + a$	$s, z z$ $s, A + a$	$s', A + s', a$	$s, A - s, a$
$z s', x$	$s', A + s', a$	$s, A + s, a$	$s', z x$ $s', A - a$	$s, z x$ $s, A - a$
r	s', z	s, z	s', x	s, x
f', z	r	t, z		
t, z	s, z	$f', z - s', z$		
f', z	t, z	r		
$t, z + f', z$	f', z	f', z		
f', x			r	t, x
t, x			s, x	$f', x - s', x$

From hence, almost an infinite number of theorems may easily be derived; some of which are the following, given here as examples of the use of the table.

I. $s, z x s, x = \frac{1}{2} r x s, a - s, A = \frac{1}{2} r x s, z - x - s, z + x = \frac{s, z}{f', x} r r = \frac{s, x}{f', z} r r.$
 $s', z x s', x = \frac{1}{2} r x s, a + s, A = \frac{1}{2} r x s, z - x + s, z + x = \frac{s', z}{f', x} r r = \frac{s', x}{f', z} r r.$

II. If A, B, C, be any three angles; $Z = A + B, X = A - B, H = \frac{1}{2} A + B + C.$
 Then $\frac{1}{2} r x w, C - w, X = s, \frac{1}{2} C + X x s, \frac{1}{2} C - X = s, \frac{1}{2} A + C - B x s, \frac{1}{2} B + C - A = s, H - B x s, H - A.$
 And $\frac{1}{2} r x w, Z - w, C = s, \frac{1}{2} Z + C x s, \frac{1}{2} Z - C = s, \frac{1}{2} A + B + C x s, \frac{1}{2} A + B - C = s, H x s, H - C.$

III. $\frac{s s, z}{s', s', z} = \frac{t t, z}{t', t', z} = \frac{r r}{r r} = \frac{v, z z}{v, z z} = \frac{t, z}{t, z};$ Or $\frac{s s, \frac{1}{2} z}{s s, \frac{1}{2} z} = \frac{t t, \frac{1}{2} z}{t t, \frac{1}{2} z} = \frac{r r}{r r} = \frac{v, z}{v, z} = \frac{t, \frac{1}{2} z}{t, \frac{1}{2} z}.$
 IV.

$$\text{IV. } \frac{1}{2}r = \frac{ss, z}{v, 2z} = \frac{ss, \frac{1}{2}z}{v, z} = \frac{s's', z}{v, 2z} = \frac{s's', \frac{1}{2}z}{v, z}; \text{ and } s, z = \frac{2ss, \frac{1}{2}z}{t, \frac{1}{2}z} = \frac{2s's', \frac{1}{2}z}{t', \frac{1}{2}z}.$$

$$\text{V. } \frac{s, z}{v, z} = \frac{r}{t, \frac{1}{2}z} = \frac{t', \frac{1}{2}z}{r} = \frac{v', z}{s, z}.$$

$$\text{VI. } \frac{t, z}{t, x} = \frac{s, A + s, a}{s, A - s, a} = \frac{t', x}{t', z}; \text{ and } \frac{rr}{t, z \times t, x} = \frac{t', z}{t, x} = \frac{t', x}{t, z} = \frac{s', a + s', A}{s', a - s', A} = \frac{t', z \times t', x}{rr}.$$

$$\text{VII. } \frac{s, A}{s, a} = \frac{t, z + t, x}{t, z - t, x} = \frac{\overline{s, z + x}}{s, z - x}; \text{ if } z \text{ and } x \text{ are two arcs, then } A = z + x, a = z - x.$$

$$\text{VIII. } \overline{s, z \pm x} = \frac{s, z \times s', x \pm s', z \times s, x}{r} = \frac{t, z \pm t, x}{f, z \times f, x}.$$

$$\text{IX. } s', \overline{z \pm x} = \frac{s', z \times s', x \pm s, z \times s, x}{r} = \frac{rr \pm t, z \times t, x}{f, z \times f, x} r.$$

$$\text{X. } \overline{t, z \pm x} = \frac{t, z \pm t, x}{rr \pm t, z \times t, x} rr; \text{ and } t', \overline{z \pm x} = \frac{rr \pm t, z \times t, x}{t, z \pm t, x}.$$

$$\text{XI. } f, \overline{z \pm x} = \frac{f, z \times f, x}{rr \pm t, z \times t, x} r; \text{ and } f', \overline{z \pm x} = \frac{f, z \times f, x}{t, z \pm t, x}.$$

XII. In three equidifferent arcs A, z, a ; where $z (= \frac{1}{2} \overline{A + a})$ is the mean arc, and $(= \frac{1}{2} \overline{A - a})$ their common difference; put $p = \frac{s, x}{r}, q = \frac{s', x}{r}$; $P = 2p \times s, z, Q = 2q \times s', z$.

Then $s, A = P - s, a = Q + s, a$; And $s, a = P - s, A = s, A - Q$.

XIII. Let $d = v, A - v, a = s', a - s', A$; then $ss, A - ss, a = \overline{2s', A \mp d} \times d = \overline{2s', a - d} \times d$.

Notes, When an arc is terminated in the second, third, or fourth quadrant, some of the signs (+ and -) of the terms in the preceding theorems, will, by the known rules, become contrary to what they now are.

XIV.

XIV. Let $A, B, C, \&c.$ be the sines, $a, b, c, \&c.$ the co-sines, $a', b', c', \&c.$ the tangents, of the arcs, $\alpha, \beta, \gamma, \&c.$ whose number is n ; the radius being r ; put S for the product of the n co-sines, $S', S'', S''', \&c.$ for the sum of the products made of every sine, every two, three, $\&c.$ sines, by the other $(n-1, n-2, n-3, \&c.)$ co-sines, where the co-sine noted by $n-n$ is unity.

Then the sine of $\overline{\alpha+\beta+\gamma+\delta, \&c.} = S' - S'' + S''' - S''', \&c. \times \frac{1}{r^{n-1}}.$

And the co-sine of $\overline{\alpha+\beta+\gamma+\delta, \&c.} = S - S' + S'' - S''', \&c. \times \frac{1}{r^{n-1}}.$

XV. Also putting T' for the sum of the tangents of the arcs, $\alpha, \beta, \gamma, \&c.$ $T'', T''', T''', \&c.$ for the sum of the products of every two, three, four, $\&c.$ tangents; and

$$\begin{aligned} A &= T' \\ B &= AT'' - T'' \\ C &= BT''' - AT'' + T'' \\ D &= CT'''' - BT''' + AT'' - T'' \\ E &= DT'''' - CT''' + BT'' - AT'' + T'' \quad \text{Put } R = \frac{1}{rr} \\ \&c. \end{aligned}$$

Then the tangent of $\overline{\alpha+\beta+\gamma+\delta, \&c.} = A + BR + CR^2 + DR^3 + ER^4; \&c.$

XVI. Hence, the sine, tangent, and secant, of any arc a , being represented by s, t, f , the co-sine, co-tangent, and co-secant, by s', t', f' ; those of the arc na are expressed as in the following theorems.

Putting $n' = n, \frac{n-1}{2}; n'' = n', \frac{n-2}{3}; n''' = n'', \frac{n-3}{4}; n'''' = n''', \frac{n-4}{5}; \&c.$

Sine of $na = \overline{nA - n''AP + n''BP - n'''CP + n''''DP, \&c.} \times \frac{s^{n-1}}{r^{n-1}};$

where $P = \frac{ss'}{r}$; $A = s; B = AP; C = BP; D = CP; \&c.$

Or $= \overline{ns - \frac{n-1}{2} \cdot \frac{n-2}{3} AP + \frac{n-3}{4} \cdot \frac{n-4}{5} BP - \frac{n-5}{6} \cdot \frac{n-6}{7} CP, \&c.} \times \frac{s^{n-1}}{r^{n-1}};$

where $A, B, C, \&c.$ stand for the respective preceding terms.

Or $= ns + \frac{1+n}{2} \cdot \frac{1-n}{3} A\mathcal{Q} + \frac{3+n}{4} \cdot \frac{3-n}{5} B\mathcal{Q} + \frac{5+n}{6} \cdot \frac{5-n}{7} C\mathcal{Q} + \frac{7+n}{8} \cdot \frac{7-n}{9} D\mathcal{Q}, \&c.$

where $\mathcal{Q} = \frac{ss'}{rr}$; $A, B, C, \&c.$ stand as before.

XVII. Co-sine of $na = \frac{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4} \times \frac{s^n}{r^{n-1}}$, where $P = \frac{t}{r}$

$$\text{Or} = r + \frac{0 - \frac{1}{2} n}{1} A \mathcal{Q} + \frac{2 - \frac{1}{2} n}{3} \frac{2 - n}{4} B \mathcal{Q} + \frac{4 - \frac{1}{2} n}{5} \frac{4 - n}{6} C \mathcal{Q} + \frac{6 - \frac{1}{2} n}{7} \frac{6 - n}{8} D \mathcal{Q} \mathcal{E} c.$$

where $\mathcal{Q} = \frac{ss}{r}$; and $A, B, C, \mathcal{E} c.$ stand for the respective preceding terms.

$$\text{Or put } M = \frac{2s^n}{r} \times r; N = \frac{rr}{4s^3}; A = \frac{1}{2}; B = AN; C = BN; D = CN, \mathcal{E} c; p = n; p' = n - 1; p'' = n - 2, \mathcal{E} c.$$

$$\text{And } a' = p; b' = p'; p'' = p; \frac{1}{2} p''; \frac{1}{3} p''; d' = p; \frac{1}{2} p''; \frac{1}{3} p''; \frac{1}{4} p''; e' = p; \frac{1}{2} p''; \frac{1}{3} p''; \frac{1}{4} p''; \frac{1}{5} p''; \mathcal{E} c.$$

$$\text{The co-sine of } na = \frac{A - Ba' + Cb' - Dc' + Ed', \mathcal{E} c. \times M.}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4}$$

XVIII. Let $A = -n' + nn'$	$A' = \frac{1}{n} n'' - n'$
$B = \frac{1}{2} n'' - nn'' + An'$	$B' = \frac{1}{n} n'' A' - n'' + n'''$
$C = -n'' + nn'' + Bn' - An'''$	$C' = \frac{1}{n} n'' B' - n'' A' + n''' - n''$
$D = \frac{1}{2} n''' - nn''' + Cn' - Bn''' + An''$	$D' = \frac{1}{n} n'' C' - n'' B' + n'' A' - n''' + n''$
$\mathcal{E} c.$	$\mathcal{E} c.$

$$\text{The tangent of } na = nt + At^3 r^{-2} + Bt^5 r^{-4} + Ct^7 r^{-6} + Dt^9 r^{-8} \mathcal{E} c.$$

$$\text{Or} = \frac{n + AN + BN^2 + CN^3 + DN^4 + \mathcal{E} c. \times t}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4}, \text{ where } N = \frac{t}{r}$$

$$\text{Or} = na' + Ab' + Bc' + Cd' + De', \mathcal{E} c. \text{ where } a' = t; b' = Na'; c' = Nb'; d' = Ne'; \mathcal{E} c.$$

$$\text{Or} = \frac{n - n' N + n'' N^2 - n''' N^3 + n^{(4)} N^4, \mathcal{E} c. \times t}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4} \times t.$$

$$\text{Co-tang}^t \text{ of } na = \frac{r^2 + A't^2 + B't^2 N + C't^2 N^2 + D't^2 N^3 + E't^2 N^4 \mathcal{E} c. \times \frac{rr}{nt}}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4}, \text{ where } N = \frac{t}{r}$$

$$\text{Or} = \frac{1 + A'N + B'N^2 + C'N^3 + D'N^4 + E'N^5, \mathcal{E} c. \times \frac{1}{t}}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4}, \text{ where } N = \frac{t}{r}$$

$$\text{Or} = \frac{1 - n' N + n'' N^2 - n''' N^3 + n^{(4)} N^4 - n^{(5)} N^5, \mathcal{E} c. \times \frac{rr}{t}}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4}, \text{ where } N = \frac{t}{r}$$

XIX. Let $A = n'$	$A' = \frac{1}{n} n''$
$B = An' - n''$	$B' = \frac{1}{n} n'' A' - n''$
$C = Bn' - An''' + n''$	$C' = \frac{1}{n} n'' B' - n'' A' + n''$
$D = Cn' - Bn''' + An'' - n'''$	$D' = \frac{1}{n} n'' C' - n'' B' + n'' A' - n'''$
$\mathcal{E} c.$	$\mathcal{E} c.$

$$\text{Secant of } na = \frac{1 + AN + BN^2 + CN^3 + DN^4 + EN^5, \mathcal{E} c. \times M}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4}$$

$$\text{Or} = \frac{1}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4} \times M; \text{ where } N = \frac{t}{r}, M = \frac{r^n}{r^n}$$

$$\text{Co-secant of } na = \frac{1 + A'N + B'N^2 + C'N^3 + D'N^4 + E'N^5, \mathcal{E} c. \times M}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4}, \text{ where } N = \frac{t}{r}, M = \frac{rr^n}{nr^n}$$

$$\text{Or} = \frac{1}{1 - n^2 P + n^{2''} P^2 - n^{4''} P^3 + n^{6''} P^4} \times M; \text{ where } N = \frac{t}{r}, M = \frac{r^{n-2}}{t^{1-2}}$$

XX. Let c be the chord of an arc (a) of the circumference of a circle, whose diameter is d . Put $N = \frac{cc}{da}$.

The chord of $na = nc + \frac{1+n}{2} \cdot \frac{1-n}{3} AN + \frac{3+n}{4} \cdot \frac{3-n}{5} BN + \frac{5+n}{6} \cdot \frac{5-n}{7} CN + \frac{7+n}{8} \cdot \frac{7-n}{9} DN, \&c.$
where $A, B, C, \&c.$ stand for the respective preceding terms.

As the preceding theorems are easily deduced from the first, the following are most readily seen to be the immediate consequences of these; and all depending upon no other principles than what are generally made use of in common computations.

XXI. Putting $s, s', t, t', f, f',$ for the sine, co-sine, tangent, co-tangent, secant, co-secant, of an arc (a), and v its versed sine; let $q' = \frac{1}{2}; q'' = \frac{1}{3}q'; q''' = \frac{1}{4}q''; q^{iv} = \frac{1}{5}q'''; q^v = \frac{1}{6}q^{iv}; \&c. N = \frac{aa}{rr}$.

$$\begin{aligned} \text{Then } s &= 1 - q''N + q^{iv}N^2 - q^{vi}N^3 + q^{viii}N^4 + q^xN^5, \&c. \times a. \\ &= a - q''a^3r^{-2} + q^{iv}a^5r^{-4} - q^{vi}a^7r^{-6} + q^{viii}a^9r^{-8}, \&c. \\ &= a - \frac{1}{2.3} AN + \frac{1}{4.5} BN - \frac{1}{6.7} CN + \frac{1}{8.9} DN, \&c. \text{ where } A, B, C, \&c. \end{aligned}$$

stand for the respective preceding terms.

$$\begin{aligned} \text{And } s' &= r - q'a^2r^{-1} + q''a^4r^{-3} - q^{iv}a^6r^{-5} + q^{vi}a^8r^{-7}, \&c. \\ &= 1 - q'N + q''N^2 - q^{iv}N^3 + q^{vi}N^4 - q^xN^5, \&c. \times r. \\ &= r - \frac{1}{1.2} a^2r^{-1} + \frac{1}{3.4} AN - \frac{1}{5.6} BN + \frac{1}{7.8} CN, \&c. A, B, C, \&c. \text{ as before.} \end{aligned}$$

$$\begin{aligned} \text{XXII. Also } v &= q'a^2r^{-1} - q^{iv}a^4r^{-3} + q^{vi}a^6r^{-5} - q^{viii}a^8r^{-7}, \&c. \\ &= \frac{1}{1.2} a^2r^{-1} - \frac{1}{3.4} AN + \frac{1}{5.6} BN - \frac{1}{7.8} CN - \frac{1}{9.10} DN, \&c. \\ &= \frac{1}{1.2} N - \frac{1}{3.4} AN - \frac{1}{5.6} BN - \frac{1}{7.8} CN, \&c. \times r. A, B, C, \&c. \text{ as before.} \end{aligned}$$

$$\begin{array}{ll} \text{XXIII. Let } A = +q' - q'' & \text{And } A' = -A \\ B = -q''' + q^{iv} + Aq' & B' = -B - AA' \\ C = +q^v - q^{vi} + Bq' - Aq''' & C' = -C - BA' - AB' \\ D = -q^{viii} + q^{x} + Cq' - Bq^{iv} + Aq^v & D' = -D - CA' - BB' - AC'. \\ \&c. & \&c. \end{array}$$

$$\text{Tangent } t = a + Aa^3r^{-2} + Ba^5r^{-4} + Ca^7r^{-6} + Da^9r^{-8}, \&c.$$

$$\text{Or } = 1 + AN + BN^2 + CN^3 + DN^4 + EN^5, \&c. \times a.$$

$$\text{Co-tangent } t' = a^{-1}r^3 + Aa + B'a^3r^{-2} + C'a^5r^{-4} + D'a^7r^{-6}, \&c.$$

$$\text{Or } = rr + Aa^2 + B'Na^2 + CN^2a^2 + D'N^2a^2, \&c. \times \frac{1}{a}.$$

XXIV.

XXIV. Also let $a = +q'$

And $a' = +q''$

$$\beta = -q''' + aq'$$

$$\beta' = -q'''' + a'q''$$

$$\gamma = +q'' - aq''' + \beta q'$$

$$\gamma' = +q'' - a'q'''' + \beta'q''$$

$$\delta = -q'''' + aq'' - \beta q''' + \gamma q'$$

$$\delta' = -q'''' + a'q'' - \beta'q'''' + \gamma'q''$$

$\mathcal{E}c.$

$\mathcal{E}c.$

$$\text{Secant } f = r + aa^2r^{-1} + \beta a^4r^{-3} + \gamma a^6r^{-5} + \delta a^8r^{-7}, \mathcal{E}c.$$

$$\text{Or} = \frac{1 + aN + \beta N^2 + \gamma N^3 + \delta N^4}{r}, \mathcal{E}c. \times r.$$

$$\text{Co-secant } f = a^{-1}r^2 + a'a + \beta'a^3r^{-2} + \gamma'a^5r^{-4} + \delta'a^7r^{-6}, \mathcal{E}c.$$

$$\text{Or} = \frac{rr + a'aa + \beta'Naa + \gamma'N^2aa + \delta'N^3aa}{r}, \mathcal{E}c. \times \frac{1}{r}. \text{ where } N = \frac{a}{r}$$

$$\text{XXV. Putting } p' = \frac{1}{2}; p'' = \frac{3}{4}p; p''' = \frac{5}{6}p''; p'''' = \frac{7}{8}p'''; p'''' = \frac{9}{10}p''''; \mathcal{E}c. N = \frac{s}{r}$$

$$\text{Then arc } a = 1 + \frac{1}{3}p'N + \frac{1}{5}p''N^2 + \frac{1}{7}p'''N^3 + \frac{1}{9}p''''N^4, \mathcal{E}c. \times s.$$

$$\text{Or} = s + \frac{1}{3}p'AN + \frac{1}{5}p''BN + \frac{1}{7}p'''CN + \frac{1}{9}p''''DN, \mathcal{E}c.$$

$$\text{Or} = s + \frac{1.1}{2.3}AN + \frac{3.3}{4.5}BN + \frac{5.5}{6.7}CN + \frac{7.7}{8.9}DN, \mathcal{E}c. \text{ where } A, B, C, \mathcal{E}c. \text{ stand for the respective preceding terms.}$$

$$\text{XXVI. If } v \text{ is the verfed sine of an arc } a, \text{ diameter being } d, M = \frac{v}{d}, R = \sqrt{dv}.$$

$$\text{Then arc } a = 1 + \frac{1.1}{2.3}M + \frac{3.3}{4.5}AM + \frac{5.5}{6.7}BM + \frac{7.7}{8.9}CM, \mathcal{E}c. \times R; A, B, C, \mathcal{E}c. \text{ are as before.}$$

$$\text{XXVII. And putting } N = \frac{r}{r}, A = t, B = AN, C = BN, D = CN, \mathcal{E}c.$$

$$\text{Then arc } a = t - \frac{1}{3}AN + \frac{1}{5}BN - \frac{1}{7}CN + \frac{1}{9}DN + \frac{1}{11}EN, \mathcal{E}c.$$

$$\text{Or} = 1 - \frac{1}{3}N + \frac{1}{5}N^2 - \frac{1}{7}N^3 + \frac{1}{9}N^4 + \frac{1}{11}N^5, \mathcal{E}c. \times t.$$

$$\text{XXVIII. Also, if } c \text{ is the chord of an arc } (a); \text{ and } N = \frac{cc}{dd}.$$

$$\text{Then arc } a = c + \frac{1.1}{2.3}AN + \frac{3.3}{4.5}BN + \frac{5.5}{6.7}CN + \frac{7.7}{8.9}DN, \mathcal{E}c. \text{ where } A, B, C, \mathcal{E}c. \text{ stand for the respective preceding terms.}$$